



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Transform[®] WG

EPA Reg. No. 62719-625

For Control of Sugarcane Aphid (*Melanaphis sacchari*) in Forage Sorghum

Section 18 Emergency Exemption

File symbol: 14FL06

FOR DISTRIBUTION AND USE ONLY IN FLORIDA UNDER SECTION 18 EMERGENCY EXEMPTION
This Section 18 Emergency Exemption is effective September 24, 2014 and expires December 31, 2014

- This labeling must be in the possession of the user at the time of application.
- It is in violation of federal law to use this product in a manner inconsistent with its labeling.
- All application directions, restrictions, and precautions on the registered product label for Transform WG (EPA Reg. No. 62719-625) are to be followed.
- Any adverse effects resulting from the use of Transform WG under this emergency exemption must be immediately reported to the Florida Department of Agriculture.

Directions for Use

Pests and Application Rates:

Pests	Transform WG (oz/acre)	Comments
Sugarcane aphid	0.75 – 1.5 (0.023 – 0.047 lb ai/acre)	Use a higher rate in the rate range for heavy pest populations.

Application Timing: Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest for forage.
- A restricted entry interval (REI) of 24 hours must be observed.
- Do not make more than two applications per acre per year.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not apply more than a total of 3.0 oz of Transform WG (0.09 lb ai of sulfoxaflor) per acre per year.

Environmental Hazards

This product is highly toxic to bees exposed through contact during spraying and while spray droplets are still wet. This product may be toxic to bees exposed to treated foliage for up to 3 hours following application. Toxicity is reduced when spray droplets are dry.

Risk to managed bees and native pollinators from contact with pesticide spray or residues can be minimized when applications are made before 7:00 am or after 7:00 pm local time or when the temperature is below 55 degrees F at the site of application.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R396-030

Approved: 09/24/14

Initial printing.