

GLYSTAR PLUS

SUPPLEMENTAL LABELING
EPA SLN No.FL-170002

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

FOR CONTROL OF BROADLEAF WEEDS IN PERENNIAL PEANUTS GROWN FOR HAY

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt..... 41.0%

OTHER INGREDIENTS: 59.0%

TOTAL 100.0%

Group	9	Herbicide
-------	---	-----------

*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid glyphosate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in possession of user at the time of application.

USE INFORMATION

PRODUCT DESCRIPTION: This product is a post-emergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

WEED RESISTANCE MANAGEMENT

Glyphosate, the active ingredient in this product, is a Group 9 herbicide. Target site resistance to Group 9 herbicides is rare. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or using other cultural practices or mechanical practices.

Refer to the product container label for additional information and instructions on resistant management

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

- Ground Broadcast Spray – Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.
- Selective Equipment –Wiper applicators and sponge bars.

DRIFT PRECAUTION

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation. Extreme care must be exercised to avoid contact of spray with foliage, green stems or fruit of desirable crops, plants, trees or other desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was NOT intended. Examples of, but not limited to, crop types that may be sensitive to glyphosate exposure include rice, small grain cereals, peanuts, potatoes, vegetables, fruits and ornamentals.

Applicators should be aware of any potentially sensitive crops near application zone before making application. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

If unsure of appropriate buffer zone, contact your local Extension Agent for advice.

PERENNIAL PEANUTS GROWN FOR HAY		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Pre-plant	To ensure a weed free field, make applications prior to planting. Apply as a broadcast application at 16 to 64 fluid ounces per acre. MAKE APPLICATIONS ACCORDING TO THE RATES LISTED IN "ANNUAL WEEDS" and "PERENNIAL WEEDS" IN THE PRODUCT CONTAINER LABEL.	Do not apply after rhizomes have been planted.
Dormant Season	This product will control or suppress many weeds including little barley, Carolina geranium and red sorrel in dormant crop. Apply 32 to 48 fluid ounces per acre of this product as a broadcast application in a minimum of 15 gallons of water per acre. Apply only to perennial peanut that is in full dormant stage. Applications may not fully control weeds damaged or under stress by frost.	Do not make more than one dormant application per year. Do not apply to crop that exhibits signs of beginning spring growth.
In Season Wiper applications	This product may be applied with wiper applicators to control or suppress the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of the container label. Application should be made at 32 to 48 fluid ounces per acre diluted to a 50% concentration to be effective.	Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting. DO NOT make wiper applications if the weed height above the crop is insufficient to ensure no application to the growing crop.
In Season Broadcast application	If weed height difference between the weeds and growing crop are not enough to make wiper applications effective, apply a 12 to 32 fluid ounce per acre rate. Use lower rates on crop stands less than 3 years old and not before crop is one year old. Well established stands tolerate higher rates. Crop will usually recover from any injury within 40 to 60 days after broadcast applications. Use in a minimum of 15 GPA dilution rate.	Do not use this product where yield reduction at next cutting cannot be tolerated. If a second application is required wait 28 days between applications. Do not exceed 48 fluid ounces per acre for all in season applications. Do not graze or harvest crop for 56 days after in season broadcast application to growing crop.
RESTRICTION FOR PERENNIAL PEANUTS: <ul style="list-style-type: none"> • Do not apply more than 2 Quarts/Acre for Pre-plant application • Do not apply more than 1.5 Quarts/Acre for Dormant season application • Do not apply more than 1.5 Quarts/Acre for all In-Season applications • Do not apply more than 5.0 Quarts/Acre per year for all uses combined 		

EPA Reg. No. 42750-61

Manufactured By:
Albaugh, LLC
Ankeny, IA 50021