

Cool-Season Food Plot Forages				
Crop	Recommended Varieties	Seeding Rates Broadcast	Planting Depth	Drainage Adaptability
Legumes				
White Clover	Ocoee, Osceola, Regal Ladino, Durana, Regalgraze, Louisiana S-1, Barblanca	3-4 lbs./ac	0 – 1/4in.	Prefers heavier, wetter soils; Not suitable for excessively drained sandy soils
Red Clover	Southern Belle, Barduro, Red Ace, Bulldog Red	10-15 lbs./ac	1/4 - 1/2in.	Prefers heavier, well drained soils
Crimson Clover	Dixie, AU-Robin, AU-Sunrise	20-25 lbs./ac	1/4 – 1/2in.	Prefers heavier, well drained soils
Arrowleaf Clover	Apache, Blackhawk	8-12 lbs./ac	0 – 1/2in.	Prefers heavier, well drained soils
Ball Clover	Don, Grazer's Select	2-3 lbs./ac	0 – 1/4in.	Wide range of soils, except excessively wet
Berseem Clover	Bigbee, Frosty	15-20 lbs./ac	1/4 – 1/2in.	Wide range of soils, except excessively wet
Medics	Armadillo, Devine Little Bur	10-15 lbs./ac (rates differ)	0 – 1/4in.	Wide range of soils, except excessively wet
Vetch	Hairy, Cahaba White, AU-Merit	20-30 lbs./ac	1 – 2in.	Wide range of soils, except excessively wet
Winter Pea	Austrian, Maple, Whistler	40-60 lbs./ac	1/2 – 1in.	Wide range of soils, except excessively wet
Grasses				
Oat	Legend 567, Horizon 720, Horizon 306, Ram, Boss Wildlife Forage, Plot Spike LA 9339, Buck Forage, Juggernaut	100-120 lbs./ac	1 – 2in.	Prefers heavier, well drained soils
Triticale	TriCal 342, TriCal 1143	90-120 lbs./ac	1 – 2in.	Wide range of soils, except excessively wet
Wheat	AGS 2024, AGS 3026, AGS4023, AGS 4043, Pioneer 26R10, Dyna-Gro Plantation	90-120 lbs./ac	1 – 2in.	Prefers heavier, well drained soils
Brassicas				
Kale	Premier, Vates, Siberian	4-5 lbs./ac	0 – 1/2in.	Wide range of soils, except excessively wet
Rape	Rangi, Rangiora, Barnapoli, Dwarf Essex, Emerald, and Winfred	4-5 lbs./ac	0 – 1/2in.	Wide range of soils, except excessively wet
Turnip	Purple Top, White Globe, Barkant	2-3 lbs./ac	0 – 1/2in.	Wide range of soils, except excessively wet
Radish	Trophy, Daikon	4-5 lbs./ac	0 – 1/2in.	Well adapted to light, sandy soils
Chicory	Puna, Forage Feast	3-5 lbs./ac	0 – 1/2in.	Well or moderately well drained soils with good fertility
<ul style="list-style-type: none"> - Food plots should consist of a blend of several forages. - In blends, the seeding rate of each component should be approximately half of the higher amount shown in the chart. 				